

Note: Brownian motion of colloidal particles of arbitrary shape

Bogdan Cichocki

*Institute of Theoretical Physics, Faculty of Physics,
University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland*

Maria L. Ekiel-Jezewska* and Eligiusz Wajnryb

*Institute of Fundamental Technological Research,
Polish Academy of Sciences, Pawińskiego 5B, 02-106 Warsaw, Poland
(Dated: January 12, 2016)*

The analytical expressions for the time-dependent cross-correlations of the translational and rotational Brownian displacements of a particle with arbitrary shape are derived. The reference center is arbitrary, and the reference frame is such that the rotational-rotational diffusion tensor is diagonal.

In general, the diffusion and mobility matrices of a particle depend on the choice of the reference center which is used to describe the rotational and translational motion. The requirement that the rotational-translational mobility matrix $\boldsymbol{\mu}^{rt}$ is symmetric defines uniquely the so-called center of mobility. In Ref. [1], this center was used to analyze the Brownian motion of a particle with arbitrary shape, and the corresponding analytical expressions for the time-dependent cross-correlations of the translational and rotational Brownian displacements were derived.

In this note, we derive the analogical analytical expressions for the cross-correlations of the translational and rotational displacements of an *arbitrary* reference center. We denote the time-dependent position of this center as $\mathbf{R}(t)$, and we introduce three mutually perpendicular unit vectors $\mathbf{u}^{(p)}(t)$, $p = 1, 2, 3$, to describe the particle orientation at time t [1, 2]. The translational and rotational Brownian displacements are,

$$\Delta \mathbf{R}(t) = \mathbf{R}(t) - \mathbf{R}(0), \quad (1)$$

$$\Delta \mathbf{u}(t) = \frac{1}{2} \sum_{p=1}^3 \mathbf{u}^{(p)}(0) \times \mathbf{u}^{(p)}(t). \quad (2)$$

Their dynamical cross-correlations are evaluated as the

averages $\langle \dots \rangle_0$ of their products with respect to the particle positions and orientations, using the conditional probability which satisfies the Smoluchowski equation [3, 4], with the diffusion matrix,

$$\mathbf{D} = \begin{bmatrix} \mathbf{D}^{tt} & \mathbf{D}^{tr} \\ \mathbf{D}^{rt} & \mathbf{D}^{rr} \end{bmatrix}, \quad (3)$$

We adopt the frame of reference in which $\mathbf{D}^{rr}$ is diagonal, i.e. $\mathbf{D}_{\mu\nu}^{rr} = D_{\mu} \delta_{\mu\nu}$, with $\mu, \nu = 1, 2, 3$.

When an arbitrary reference center is chosen, the tensor $\mathbf{D}^{rt}$ contains non-vanishing antisymmetric part. In this case to derive the rotational-translational correlations one can proceed as it is presented in Sec. VIII of Ref. [1] with the following modification. The correlation $\langle \mathbf{u}^{(p)}(t) \Delta \mathbf{R}(t) \rangle_0$ is evaluated in the analogical way as in Eq. (79) from Ref. [1], but with another second rank tensor $\mathbf{A}^{(p)}$, i.e.

$$\mathbf{A}^{(p)} = \mathbf{u}^{(p)} \times \mathbf{D}^{rt} - \frac{1}{2} \mathbf{u}^{(p)} \mathbf{V}, \quad (4)$$

where the components of vector $\mathbf{V}$ are given as $V_{\alpha} = \sum_{\mu, \nu=1}^3 \epsilon_{\alpha\mu\nu} D_{\mu\nu}^{rt}$.

The resulting off-diagonal Cartesian components are,

$$\begin{aligned} \langle \Delta \mathbf{u}(t) \Delta \mathbf{R}(t) \rangle_{0, \alpha\beta} = & - \left[(D_{\alpha\beta}^{rt} + D_{\beta\alpha}^{rt}) \left(\frac{1}{4} + \frac{D_{\beta} - D_{\gamma}}{4\Delta} \right) + (D_{\alpha\beta}^{rt} - D_{\beta\alpha}^{rt}) \left(\frac{1}{6} + \frac{D - D_{\alpha}}{2\Delta} \right) \right] \frac{e^{-f^{(-)}t} - e^{-f_{\gamma}^{(1)}t}}{f^{(-)} - f_{\gamma}^{(1)}} \\ & - \left[(D_{\alpha\beta}^{rt} + D_{\beta\alpha}^{rt}) \left(\frac{1}{4} - \frac{D_{\beta} - D_{\gamma}}{4\Delta} \right) + (D_{\alpha\beta}^{rt} - D_{\beta\alpha}^{rt}) \left(\frac{1}{6} - \frac{D - D_{\alpha}}{2\Delta} \right) \right] \frac{e^{-f^{(+)}t} - e^{-f_{\gamma}^{(1)}t}}{f^{(+)} - f_{\gamma}^{(1)}} \\ & - \frac{D_{\alpha\beta}^{rt} + D_{\beta\alpha}^{rt}}{4} \left[\frac{e^{-f_{\alpha}^{(2)}t} - e^{-f_{\beta}^{(1)}t}}{f_{\alpha}^{(2)} - f_{\beta}^{(1)}} + \frac{e^{-f_{\alpha}^{(1)}t} - e^{-f_{\beta}^{(1)}t}}{f_{\alpha}^{(1)} - f_{\beta}^{(1)}} \right] - \frac{D_{\alpha\beta}^{rt} - D_{\beta\alpha}^{rt}}{2} \left[\frac{e^{-f_{\alpha}^{(2)}t} - e^{-f_{\beta}^{(1)}t}}{f_{\alpha}^{(2)} - f_{\beta}^{(1)}} - \frac{1 - e^{-f_{\gamma}^{(1)}t}}{3f_{\gamma}^{(1)}} \right], \end{aligned} \quad (5)$$

with the convention that everywhere in this note (α, β, γ) is a permutation of $(1, 2, 3)$, and

$$f_{\alpha}^{(1)} = 3D - D_{\alpha}, \quad f_{\alpha}^{(2)} = 3(D_{\alpha} + D), \quad \alpha = 1, 2, 3, \quad (6)$$

$$f^{(\pm)} = 6D \pm 2\Delta, \quad (7)$$

$$D = \frac{1}{3}(D_1 + D_2 + D_3), \quad (8)$$

$$\Delta = \sqrt{D_1^2 + D_2^2 + D_3^2 - D_1 D_2 - D_1 D_3 - D_2 D_3}. \quad (9)$$

The diagonal Cartesian components read:

$$\begin{aligned} \langle \Delta \mathbf{u}(t) \Delta \mathbf{R}(t) \rangle_{0,\alpha\alpha} = & -\frac{D_{\alpha\alpha}^{rt} - D_{\beta\beta}^{rt}}{8} \frac{e^{-f_{\gamma}^{(2)}t} - e^{-f_{\gamma}^{(1)}t}}{D_{\gamma}} \\ & -\frac{D_{\alpha\alpha}^{rt} - D_{\gamma\gamma}^{rt}}{8} \frac{e^{-f_{\beta}^{(2)}t} - e^{-f_{\beta}^{(1)}t}}{D_{\beta}} + \frac{D_{\alpha\alpha}^{rt} + D_{\beta\beta}^{rt}}{2} e^{-f_{\gamma}^{(1)}t} t \\ & + \frac{D_{\alpha\alpha}^{rt} + D_{\gamma\gamma}^{rt}}{2} e^{-f_{\beta}^{(1)}t} t. \end{aligned} \quad (10)$$

To derive the translational-translational correlations, the derivative (54) from Ref. [1] must be changed into

$$\begin{aligned} \frac{d}{dt} \langle \Delta \mathbf{R}(t) \Delta \mathbf{R}(t) \rangle_0 \\ = 2 \langle \mathbf{D}^{tt}(t) \rangle_0 + \langle \mathbf{V}(t) \Delta \mathbf{R}(t) \rangle_0 + \langle \Delta \mathbf{R}(t) \mathbf{V}(t) \rangle_0. \end{aligned} \quad (11)$$

The two last terms can be then calculated by decomposing the vector $\mathbf{V}$, defined under Eq. (4), in the frame $\mathbf{u}^{(p)}$, $p = 1, 2, 3$, and using the results for $\langle \mathbf{u}^{(p)}(t) \Delta \mathbf{R}(t) \rangle_0$ obtained by the method described above.

The off-diagonal Cartesian components with $\alpha \neq \beta$ read,

$$\begin{aligned} \frac{1}{2} \langle \Delta \mathbf{R}(t) \Delta \mathbf{R}(t) \rangle_{0,\alpha\beta} = & \frac{1 - e^{-f_{\gamma}^{(2)}t}}{f_{\gamma}^{(2)}} D_{\alpha\beta}^{tt} \\ & + \frac{1}{2} (D_{\beta\gamma}^{rt} - D_{\gamma\beta}^{rt}) (3D_{\gamma\alpha}^{rt} - D_{\alpha\gamma}^{rt}) I(f_{\gamma}^{(2)}, f_{\alpha}^{(1)}; t) \\ & + \frac{1}{2} (D_{\alpha\gamma}^{rt} - D_{\gamma\alpha}^{rt}) (3D_{\gamma\beta}^{rt} - D_{\beta\gamma}^{rt}) I(f_{\gamma}^{(2)}, f_{\beta}^{(1)}; t) \\ & - (D_{\alpha\beta}^{rt} - D_{\beta\alpha}^{rt}) (D_{\alpha\alpha}^{rt} - D_{\beta\beta}^{rt}) I(f_{\gamma}^{(2)}, f_{\gamma}^{(1)}; t), \end{aligned} \quad (12)$$

where

$$I(f_1, f_2; t) = \frac{1}{f_1 f_2} + \frac{e^{-f_1 t}}{f_1(f_1 - f_2)} - \frac{e^{-f_2 t}}{f_2(f_1 - f_2)}. \quad (13)$$

The diagonal Cartesian components are,

$$\begin{aligned} \frac{1}{2} \langle \Delta \mathbf{R}(t) \Delta \mathbf{R}(t) \rangle_{0,\alpha\alpha} = & W(t) + \sum_{\sigma=\pm} \frac{1 - e^{-f^{(\sigma)}t}}{f^{(\sigma)}} D_{\alpha\alpha}^{tt(\sigma)} \\ & + \sum_{\sigma=\pm} \sum_{\beta=1}^3 P^{(\sigma)}(\alpha, \beta) I(f^{(\sigma)}, f_{\beta}^{(1)}; t), \end{aligned} \quad (14)$$

where

$$\begin{aligned} D_{\alpha\alpha}^{tt(\pm)} = & \left(\frac{1}{2} \mp \frac{3}{4} \frac{D_{\alpha} - D}{\Delta} \right) \left(D_{\alpha\alpha}^{tt} - \frac{1}{3} \text{Tr} \mathbf{D}^{tt} \right) \\ & \pm \frac{D_{\beta} - D_{\gamma}}{4\Delta} (D_{\beta\beta}^{tt} - D_{\gamma\gamma}^{tt}), \end{aligned} \quad (15)$$

and

$$\begin{aligned} P^{(\pm)}(\alpha, \beta) = & -(D_{\alpha\gamma}^{rt} - D_{\gamma\alpha}^{rt})^2 \left(\frac{1}{3} \mp \frac{D - D_{\gamma}}{\Delta} \right) \\ & + [(D_{\alpha\gamma}^{rt})^2 - (D_{\gamma\alpha}^{rt})^2] \left(\frac{1}{2} \mp \frac{D_{\alpha} - D_{\beta}}{2\Delta} \right), \text{ for } \alpha \neq \beta, \end{aligned} \quad (16)$$

$$P^{(\pm)}(\alpha, \alpha) = -P^{(\pm)}(\beta, \alpha) - P^{(\pm)}(\gamma, \alpha), \quad (17)$$

Since $\sum_{\alpha=1}^3 D_{\alpha\alpha}^{tt(\pm)} = 0$, it is easy to see that $W(t)$ is 1/6 of the mean square Brownian displacement,

$$W(t) = \frac{1}{6} \langle \Delta \mathbf{R}(t) \cdot \Delta \mathbf{R}(t) \rangle_0. \quad (18)$$

Moreover, it follows from our calculation that

$$W(t) = D_{cm} t + \frac{1}{3} \sum_{\alpha=1}^3 \frac{(D_{\beta\gamma}^{rt} - D_{\gamma\beta}^{rt})^2}{(D_{\beta} + D_{\gamma})^2} (1 - e^{-(D_{\beta} + D_{\gamma})t}), \quad (19)$$

where $\beta, \gamma \neq \alpha$ and D_{cm} is the self-diffusion coefficient referring to the center of mobility,

$$D_{cm} = \frac{1}{3} \left[\text{Tr} \mathbf{D}^{tt} - \sum_{\alpha=1}^3 \frac{(D_{\beta\gamma}^{rt} - D_{\gamma\beta}^{rt})^2}{D_{\beta} + D_{\gamma}} \right]. \quad (20)$$

The importance of the center of mobility follows from Eq. (19) – this is the only reference center for which the mean square displacement is a linear function of time.[5]

For completeness, we remind the rotational-rotational correlations [1], which do not depend on the choice of the reference center. The off-diagonal elements vanish, and the diagonal elements, $\alpha = 1, 2, 3$, are given by

$$\begin{aligned} \langle \Delta \mathbf{u}(t) \Delta \mathbf{u}(t) \rangle_{0,\alpha\alpha} = & \frac{1}{6} - \frac{3(D - D_{\alpha}) + \Delta}{12\Delta} e^{-f^{(-)}t} \\ & - \frac{3(D_{\alpha} - D) + \Delta}{12\Delta} e^{-f^{(+)}t} - \frac{1}{4} e^{-f_{\alpha}^{(2)}t} + \frac{1}{4} e^{-f_{\alpha}^{(1)}t}. \end{aligned} \quad (21)$$

In this work, the analytical expressions for the cross-correlations are more complex than their analogs in Ref. [1], but the advantage is that they apply to an *arbitrary* reference center. Therefore, they can be easily used to account for the experimental results, obtained for any reference center. An example of the comparison of the theoretical expressions derived here with the measurements (made in Ref. [2]) can be found in Ref. [6].

M.L.E.-J. and E.W. were supported in part by the Polish National Science Centre under Grant No. 2012/05/B/ST8/03010. M.L.E.-J. benefited from the scientific activities of the COST Action MP1305.

* Corresponding author: mekiel@ippt.pan.pl

- [1] B. Cichocki, M. L. Ekiel-Jezewska and E. Wajnryb, J. Chem. Phys. **142**, 214902 (2015).
- [2] D. J. Kraft, R. Wittkowski, B. ten Hagen, K. V. Edmond, D. J. Pine and H. Löwen, Phys. Rev. E **88**, 050301 (2013).
- [3] N. van Kampen, *Stochastic Processes in Physics and Chemistry, 3rd Edition*, North-Holland, 2007.
- [4] R. B. Jones and P. N. Pusey, Annu. Rev. Phys. Chem. **42**, 137 (1991).
- [5] B. Cichocki, M. L. Ekiel-Jezewska and E. Wajnryb, J. Chem. Phys. **136**, 071102 (2012).
- [6] B. Cichocki, M. L. Ekiel-Jezewska and E. Wajnryb, to be published.